



## Food applications of *Irvingia gabonensis* (Aubry-Lecomte ex. O'Rorke) Baill., the 'bush mango': A review

L. Mateus-Reguengo, L. Barbosa-Pereira, W. Rembangouet, M. Bertolino, M. Giordano, O. Rojo-Poveda & G. Zeppa

To cite this article: L. Mateus-Reguengo, L. Barbosa-Pereira, W. Rembangouet, M. Bertolino, M. Giordano, O. Rojo-Poveda & G. Zeppa (2020) Food applications of *Irvingia gabonensis* (Aubry-Lecomte ex. O'Rorke) Baill., the 'bush mango': A review, Critical Reviews in Food Science and Nutrition, 60:14, 2446-2459, DOI: [10.1080/10408398.2019.1646704](https://doi.org/10.1080/10408398.2019.1646704)

To link to this article: <https://doi.org/10.1080/10408398.2019.1646704>



Published online: 01 Aug 2019.



Submit your article to this journal [↗](#)



Article views: 233



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

REVIEW



## Food applications of *Irvingia gabonensis* (Aubry-Lecomte ex. O'Rorke) Baill., the 'bush mango': A review

L. Mateus-Reguengo<sup>a</sup>, L. Barbosa-Pereira<sup>a</sup>, W. Rembangouet<sup>b</sup>, M. Bertolino<sup>a</sup>, M. Giordano<sup>a</sup>, O. Rojo-Poveda<sup>a</sup>, and G. Zeppa<sup>a</sup>

<sup>a</sup>Department of Agriculture, Forest, and Food Sciences (DISAFA), University of Turin, Turin, TO, Italy; <sup>b</sup>University of Gastronomic Sciences, Bra, CN, Italy

### ABSTRACT

*Irvingia gabonensis*, also known as 'bush mango', is a multipurpose fruit tree, native to tropical Africa. It is a priority indigenous fruit tree in western and central Africa since its wood is used for making utensils and fruits are mostly used as food and medicine. The objective of this work is to provide an updated review of the available knowledge about physicochemical characteristics of *I. gabonensis* fruit in order to evaluate its potential use in the food industry. The fruit mesocarp contains various phytochemicals and ascorbic acid concentration higher than some vitamin C rich fruits, then it is consumed fresh or dried, used to produce juice and wine, or as a flavourant. *I. gabonensis* fruit kernel is rich in oil (63%–69% crude fat), mainly composed of myristic and lauric acids. Its triacylglycerol composition and, resultantly, melting curve and polymorphism indicate an aptitude for diverse applications, as it is solid at room temperature. Forty-one phenolic compounds were identified in the seeds and derived extracts and supplements, being ellagic acid and its derivatives the most present. This review enhances our knowledge about nutritional content and health benefits of *I. gabonensis* whole fruit, especially its pulp and seed, evidencing the need for safer and more efficient production of value-added products.

### KEYWORDS

Bush mango; ogbono soup; dika nut; polyphenols; carotenoids

### Introduction

*Irvingia gabonensis* (Aubry-Lecomte ex O'Rorke) Baill., also known as 'African mango tree', 'bush mango', 'sweet bush mango', 'dika nut', 'rainy season bush mango', 'dika bread tree', 'odika', 'manguier sauvage', 'chocolatier', or 'ogbono', is a multipurpose fruit tree native to tropical Africa, and more specifically to West and Central Africa (National Research Council 2006; Singh 2007). This traditional tree is common in dense evergreen rain forests but is also found near riverbanks (Atangana et al. 2001); it has been reported to be used as a source of timber and to make utensils, and also as food and medicine (Okoronkwo et al. 2014; Fungo et al. 2016; Ofundem et al. 2017).

The fruits are available from May to September with the peak harvesting period being June/July (Onimawo et al. 2003). The fruit is a broad, ellipsoid drupe with a thin epicarp, an edible fleshy mesocarp (pulp) (when ripe) and a hard, stony, nut encasing a soft, oil rich, dicotyledonous kernel wrapped inside a brown seed-coat (Ogunsina, Koya, and Adeosun 2008a, 2008b; Ogunsina et al. 2012; Ogunsina, Olatunde, and Adeleye 2014). This kernel, which is also referred to as seed, is widely used as food (Omoniyi et al. 2017). Thus, *I. gabonensis* is included in the FAO/INFOODS biodiversity list and is recognized by the International

Centre for Research in Agroforestry as a priority indigenous fruit tree for West and Central Africa (Franzel, Jaenicke, and Janssen 1996; Leakey et al. 2005; National Research Council 2006; Ruth Charrondière et al. 2013; Stadlmayr et al. 2013; Vihotogbé, van den Berg, and Sosef 2013; Bvenura and Sivakumar 2017; Shaheen, Ahmad, and Haroon 2017). Over the last 25 years work in Cameroon and Nigeria has been also focused on the domestication of *I. gabonensis* to diversify cropping systems, generate income for subsistence farmers and to create a new paradigm for tropical agriculture based on the sustainable intensification of smallholder farming (Leakey 2014, 2018, 2019; Leakey et al. 2012). Due to this, ethnobotanical and economical researches focused on *I. gabonensis*, have emerged in recent years (Ladipo, Fondoun, and Ganga 1996; Ayuk et al. 1999; Leakey et al. 2000; Atangana et al. 2001; Leakey and Page 2006; Singh 2007; Vihotogbé, van den Berg, and Sosef 2013); however, up until now, a comprehensive review summarizing its applications and the functional properties of *I. gabonensis* products in the food industry is lacking. The aim of this review is thus to deepen the existing knowledge about the physicochemical characteristics of *I. gabonensis* pulp and kernels in order to verify their potential use in the food industry, especially in functional products.

## Literature search

Literature published prior to 2019 were examined using various databases, such as ScienceDirect, Scopus, Scielo, Google Scholars, Research Gate, PubMed and Web of Science. As *I. gabonensis* presents a specific nomenclature in each region of Africa, it was necessary to use several terms for a complete search. 'Irvingia', 'Irvingia gabonensis', '*I. gabonensis*', 'African Mango', 'Bush Mango', 'Ogbono', 'Dika Fruit', 'Dika Nut', 'Dikanut', 'Dika Kernel', 'Dika Seed' and 'Ogbono Seed' were the keywords used in the databases. Articles were also identified by a manual search of bibliographies from all the retrieved articles.

## Pulp

The fresh fruits, that are similar to small mangoes, have a green-yellow color and, since their taste varies between sweet and bitter, they are divided into two groups, the 'eating type' and the 'cooking type'. The 'eating type' comprises of the species *Irvingia gabonensis* which is fibrous, has a mesocarp characterized by a sweet taste and a yellow to orange color while the 'cooking type' is the *Irvingia wombolu* species whose seeds are widely processed across West Africa but the mesocarp is bitter and non-edible (Harris 1996).

## Chemical composition

Nutritional and chemical composition of *I. gabonensis* fruit (pulp) are shown on Table 1. The pulp of *I. gabonensis* has an elevated moisture content, from 78.8% to 90.5%, and a soluble solid content of around 10%, which indicates that this fruit is suitable for juice production. The pH varies between 4.7 and 6.2 and acidity may be the reason behind the bitter taste of the pulp (Onimawo et al. 2003). The ash content is low (0.8–1.8%) but potassium (1114 mg/100 g dry

weight) and calcium (118 mg/100 g dry weight) contents are high in contrast to low sodium content (12 mg/100 g dry weight) (Olayiwola et al. 2013). The high variability on reported fat content of this fruit could be due to the differences in sample extraction amongst different studies.

Polyphenol composition is described in the following section, as it is related to *I. gabonensis* pulp functional properties.

## Functional properties

*I. gabonensis* fruits are well cited as anti-sickling products then useful for the treatment of sickle cell disorder or sickle cell anemia, an autosomal recessive genetic blood disorder with over-dominance characterized by red blood cells with abnormal, rigid, sickle shaped that afflicting the population living in Africa, South America and Asia (Amujoyegbe et al. 2016). Etebu (2012) compared the phytochemicals in *I. gabonensis* and *I. wombolu* documenting the presence of five groups of phytochemicals (alkaloids, flavonoids, saponins, tannins and glucosides) in mesocarp from both varieties. This finding is supported by other studies which investigated these components or other (Table 2). The fruit of *I. gabonensis* can be considered vitamin C rich (51–76 mg/100 g) when compared with other fruits like orange (about 50 mg/100 g), and common mango (*Mangifera indica* L.) (about 40 mg/100 g) (USDA 2018). Also, the carotenoid and the phenolic content of *I. gabonensis* fruits are very high (Table 2).

Emejulu et al. (2014) studied the effect of *I. gabonensis* fruit juice on serum lipid profile of sodium fluoride-intoxicated rats by comparing with positive and negative control groups. They concluded that the level of HDL-cholesterol was higher in the *I. gabonensis* group than in the positive control group (20 mg/kg body weight of quercetin + 100 mg/kg body weight of alpha-tocopherol) and the fruit juice of *I. gabonensis* was reported to have a lowering effect

**Table 1.** Proximate composition of *Irvingia gabonensis* ripe raw pulp.

	FAO 1968	Onimawo et al. 2003	Stadlmayr et al. 2013	Aina 1990	Joseph and Aworh 1991	Mbaeyi and Anyanwu 2010
Moisture (%)	81.4	80.0	78.8	80.5	80.0	90.5
Ash (%)	1.8	0.8	0.8	—	—	1.4
Fiber (%)	0.4	0.4	0.4	—	—	4.9
Fat (%)	0.2	1.1	1.1	—	—	2.2
Protein (%)	0.9	1.1	1.1	—	—	4.4
Carbohydrate (%)	15.7	16.7	17.8	—	—	4.6
Total Acidity (%)	—	0.1	—	0.3	0.1	—
pH	—	5.8	—	4.7	5.0	6.2
Soluble solids (%)	—	10.0	—	10.5	14.0	9.5

**Table 2.** Phytochemical content (mg/100 g fresh fruit) of *I. gabonensis* ripe raw pulp.

		References
Ascorbic and dehydroascorbic acid	51.00–76.07	FAO 1968; Onimawo et al. 2003; Stadlmayr et al. 2013; Aina 1990; Joseph and Aworh 1991; Olayiwola et al. 2013; Achinewhu 1983
Carotenoids	1.26–2.21	Olayiwola et al. 2013; Aina 1990
Tannin	54.9	Aina 1990
Phenolics	382.20	Olayiwola et al. 2013
Vitamin A	280.18*	Mbaeyi and Anyanwu 2010

\*Retinol equivalent.

on LDL-cholesterol as compared to the other groups tested. The author attributed this action to the presence of alkaloids, saponins, flavonoids and polyphenols commonly known to reduce serum lipids in animals (Ezekwe and Obidoa 2001). An ameliorative effect was observed in NaF-induced lipidemia in rats when fed with *I. gabonensis* fruit juice, which may be due to its reportedly rich vitamin C content and plant polyphenols. *I. gabonensis* pulp is also used for diabetes treatment when coupled with *Ouratea turnarea* and *Citrus medica* (Kuethe and Efferth 2011).

### Traditional processing and products obtained

As *I. gabonensis* kernels are more economically resourceful, the traditional post-harvesting operations aim to remove the kernel in its optimum conditions. In most cases, the interest in the seed results in the neglect of the potential of other parts of the fruit, including the pulp.

Fruit harvesting must be undertaken at an appropriate time, preventing the harvest of immature fruits, but also ensuring a good shelf life (Ladipo 1999). Besides harvesting, the gathering of fallen ripe has also been reported (Elah 2010; Nkwatoh et al. 2010). The fresh fruits of *I. gabonensis* have a shelf life of less than 2 days if picked when ripe and not more than 10 days if harvested at the mature green stage, due to high respiration rate, moisture loss and microbial attack (Aina 1990; Joseph and Aworh 1991, 1992; Etebu 2012). Etebu (2012) isolated four genera of fungi (*Aspergillus*, *Penicillium*, *Rhizopus* and *Mucor*) from post-harvest fruits of *I. gabonensis* and *I. wombolu*, concluding that *Rhizopus* and *Mucor* species were the most predominant genera of fungi associated with postharvest *Irvingia* fruits.

Aina (1990) described the physicochemical changes in *I. gabonensis* fruits during normal storage ripening and revealed that, with the ripening process, the fruit peel gets yellow and the sweetness of the pulp increases due to starch degradation. The sourness and the acidity decreases, the fruit turns softer, mainly due to pectin degradation, and, finally, the vitamin C content decreases as ascorbic acid is very susceptible to oxidative degradation. In order to extend the shelf-life of *I. gabonensis* fruits during storage, Joseph and Aworh (1991, 1992) studied the influence of some post-harvest treatments. Firstly, while comparing ripening at room temperature and refrigerated storage of the fruits, it was noticed that low temperatures induced cold injuries in *I. gabonensis* and that room-ripened fruits had better flesh color and texture, although, they also had a higher moisture loss (Joseph and Aworh 1991). After these primary results, the authors conducted other experiments in order to determine the effects of dipping fruits in hot water and in different concentrations of benomyl, DHA-S, and  $\text{Na}_2\text{S}_2\text{O}_5$  at different temperatures, on the shelf life and quality. While untreated fruits had become brownish black and unmarketable by day 12, the fruits treated with hot 0.1% benomyl or 0.5%  $\text{Na}_2\text{S}_2\text{O}_5$  solutions followed by waxing had an attractive appearance and good quality until day 14. Dipping fruits in hot water (55 °C) or chemical solutions (0.1% benomyl,

0.5% DHA-S or 0–5%  $\text{Na}_2\text{S}_2\text{O}_5$ ) followed by waxing or packaging in boxes overwrapped with stretch PVC film, delayed ripening, controlled decay, minimized weight loss and extended the shelf life of the fruits under tropical ambient conditions, without adverse effects on visual and chemical qualities (Joseph and Aworh 1992).

*I. gabonensis* pulp is consumed to a considerable extent, normally eaten raw as a dessert or snack; however, large quantities are usually wasted (Akubor 1996). Juice, beverage, and jam manufacturing requires little processing and addresses the need to use the raw pulp, injured or not. Various authors cited bush mango as being more suitable for juice, wine, and jam production, compared to other known tropical fruits (Ejiofor 1994; Okolo 1994; Agbor 1994; Okafor, Okolo, and Ejiofor 1996; Akubor 1996; Ainge and Brown 2001; Aworh 2015).

Laboratory trials have shown that jam can be produced from lesser known Nigerian fruits including *I. gabonensis* (Aina and Adesina 1991; Ainge and Brown 2001; Aworh 2014, 2015). Aworh (2014) produced jams from three indigenous fruits containing 50% of pulp. For the *I. gabonensis* jam recipe, 500 g of pulp was mashed and boiled with 638 g of sugar, 100 g of water, 6 g of citric acid and 5 g of calcium chloride in a steam-jacketed kettle. In a sensory evaluation of the wild fruit jams, *I. gabonensis* jam was the less preferred, especially in terms of flavor and consistency. The author concluded that although *I. gabonensis* jam is manufacturable, it may not be marketable for its low acceptance.

Akubor (1996) studied the suitability of *I. gabonensis* fruits for juice and wine production. The pulp was blended with water in a 1:5 proportion, then filtered in cheesecloth and cane sugar added in order to obtain 23°Brix. A yield of 75% was achieved and the obtained juice was compared to other tropical fruit juices obtained from banana, orange and cashew. *I. gabonensis* juice showed a lower protein content and a higher ascorbic acid content compared to the other tropical fruit juices.

For wine production, the *I. gabonensis* juice was fermented with *Saccharomyces cerevisiae* at 30 °C for 28 days. The wine produced had 8.12% (v/v) alcohol, 0.78% protein, 6.5°Brix SS, and a pH 3.10. Consumer test showed that the obtained product was generally accepted and had no significant differences in color, sweetness, mouthfeel, and general acceptability as compared to a German reference wine.

Besides beverage production, osmotic dehydration has also been cited as an excellent application of *I. gabonensis* fruits (Falade and Aworh 2004; Falade and Aworh 2005; Aworh 2015). With this process, a variety of new shelf-stable food products can be developed with few modifications in the fruits' color, flavor, and texture characteristics. Osmose-dried products could reduce perishable fruit losses postharvest and ensure that seasonal fruit products are available throughout the year. The osmotic process is very suitable as a pretreatment prior to air-drying of fruits, resulting in a fruit product with an intermediate moisture content (Falade and Aworh 2004, 2005; Aworh 2015).

Falade and Aworh (2004) studied the influence of osmotic pretreatment on the adsorption isotherms of osmo-

air dried *I. gabonensis* fruits. The treatments were performed at 27 °C and 40 °C, with sugar concentrations of 52°Brix, 60°Brix and 68°Brix, maintaining a fruit:sirup ratio of 1:4 w/w for 10 h. Afterwards the fruit slices were oven air-dried at 60 °C for 72 h. The authors concluded that the adsorption isotherms of osmo-oven dried fruits followed a type III isotherm, which characterizes high sugar products like many other fruits. In the experiments, *I. gabonensis* isotherm was affected by fruit ripeness degree. In fact, the equilibrium moisture content of the fruit increased with higher degree of ripeness at the same water activity ( $a_w$ ), concentration of the sucrose solution used for pretreatment (the higher the sugar concentration, less water was absorbed at low and intermediate  $a_w$  ranges), and equilibrium temperature (equilibrium moisture content decreased with increasing temperature, when  $a_w < 0.8$ ).

Dried fruits of *I. gabonensis* can also be used as a flavoring agent in other food products in order to diversify its usage. Mbaeyi and Anyanwu (2010) evaluated the use of these products as a yogurt flavourant. The dried and pliable fruits were milled, sieved through a 0.59 mm sieve, and added at 0.8%, 1.6%, 2.4%, 3.2%, 4.0%, and 4.8% in commercial full-fat cow yogurt. The best sensory results were obtained in the yogurt containing 0.8% of dried fruit with an overall acceptability statistically not different from commercial yogurt.

## Seeds

The kernels are the main products of *I. gabonensis* and constitute an important part of West and Central Africa diet, mainly in rural communities, providing carbohydrate and protein. The seed consists of a hard shell, an outer brown testa (hull) and inside, the kernel, composed of two white cotyledons. The seeds of the fruits of *I. gabonensis*, can be eaten raw or roasted and are used in food preparations (National Research Council 2006).

### Chemical composition

The summary of the proximate composition of *I. gabonensis* kernels is provided in Table 3. According to literature, *I. gabonensis* seed has a high energetic value (595–729 kcal), due to high percentages of fat (10–71%), carbohydrates (3–52%) and protein (7–22%). With crude protein content ranging from 7 to 22%, *I. gabonensis* seeds have comparable or higher protein levels than the majority of the cereals comprising our daily diet (corn, sorghum, rice, etc), which generally does not exceed 13%. Fiber content was generally low in the studies reported, except by Onimawo et al. (2003), who observed an outlying value of 10.23%. This may be due to sample preparation but also the intraspecific variability that offers horticulturalists and agroforesters the opportunity to develop cultivars appropriate to different uses and industries (Leakey et al. 2012; Leakey 2017, 2019).

**Table 3.** Proximate composition of *Ivingia gabonensis* kernel.

	Reference														
	Idowu et al. 2013	Onyike and Acheru 2002	Eiah 2010	Okoronkwo et al. 2014	Ibezim 2015	Matos et al. 2009	Ekpe et al. 2007	Ogunsina et al. 2012	Joseph 1995	Onimawo et al. 2003	Dosumu et al. 2012	Ezeabara and Ezeani 2016	Giami et al. 1994	Eka 1980	Ejiofor et al. 1987
Energy (kcal)	684.5	688.00	—	—	—	729.36	—	—	—	—	595.05	—	682	—	—
Moisture content (%)	4.26	5.93	2.55	3.21	8.28	0.02	5.20	2.55	—	3.36	3.75	11.54	4.1	2.10	11.90
Protein (%)	8.10	8.71	8.90	7.47	8.81	8.52	7.60	8.90	8.70	7.76	21.52	17.43	10.9	8.60	9.24
Fat (%)	67.69	62.80	68.37	68.81	69.34	68.24	66.60	68.37	66.50	65.46	55.09	9.81	64.2	70.80	51.32
Fiber (%)	5.88	—	—	4.38	1.25	—	1.90	—	—	10.23	—	4.18	3.4	1.40	0.86
Ash (%)	3.35	0.63	2.32	8.71	4.52	2.93	9.50	2.32	—	2.26	16.35	4.66	2.2	6.80	2.46
Carbohydrate (%)	10.72	21.93	18.67	11.39	11.39	20.28	—	18.67	—	10.93	3.29	52.40	15.2	14.10	26.02



### Kernel oil composition and physical properties

The analysis revealed that *I. gabonensis* is essentially a rich source of edible fat, with a mean percentage of 62%. Some oil-bearing products with such high percentage of oil are coconut, almond, pistachio, sunflower, walnut, and watermelon seeds which contain 62.3, 58.9, 53.5, 52.1, 64.5 and 52.6% of oil, respectively (Gopalan, Rama Sari, and Balasubramanian 2014).

Physicochemical characteristics of *I. gabonensis* kernel oil are summarized in Table 4. This oilseed could also be a source of minerals with nutritional value such as iron, copper and zinc and therefore recommended for use in diets. The content in micronutrients such as trace elements (iron, copper and zinc) and microelements (potassium and sodium) of *I. gabonensis* kernel oil are summarized in Table 5. Acid value is used as an indicator for edibility of oil and suitability for use in the paint industry (Etong, Mustapha, and Taleat 2014). The acid value for samples of *I. gabonensis* oil found in literature (3.18–24.7 mg KOH/g) were considerably diverse, and most of them did not fall within the allowable limits for edible oils, 4.0 mg KOH/g fat for oil (Codex Alimentarius 2015). The free fatty acid values ranged from 0.30 to 4.70%, which can be considered low if compared with other vegetable oils (Omoniyi et al. 2017). Etong, Mustapha, and Taleat (2014) reported that a low acid value with a correspondingly low level of free fatty acid, suggests the low level of hydrolytic and lipolytic activities in the oil, thus the seed oil studied could be a good source of raw materials for industries. The peroxide value (0.04–3.33 meq O<sub>2</sub>/kg fat) was incredibly low, which indicates a low level of oxidative rancidity and also suggests a high antioxidant level in the oil (Etong, Mustapha, and Taleat 2014). Etong, Mustapha, and Taleat (2014) also stated that the relative low iodine number of the seed oil may be indicative of the presence of few unsaturated bonds and low susceptibility to oxidative rancidity. High saponification value (187.90–701.00 mg KOH/g) also indicates it has potential for industrial use (Omoniyi et al. 2017). Low unsaponifiable

matter (0.12–1.70%) indicates that the oil is pure (Etong, Mustapha, and Taleat 2014).

*I. gabonensis* seed kernel oil (Table 6) is mostly cited as a mystiric-lauric oil, with mystiric acid being the most abundant followed by lauric acid (Matos et al. 2009; Silou et al. 2011; Yamoneka et al. 2015; Lieb et al. 2018). According to Matos et al. (2009), *I. gabonensis* kernel oil is a technical fat because it resists thermo oxidative, hydrolytic, and enzymatic activities due to its fatty acid profile.

Nine free fatty acids were described in literature, only three of which are unsaturated. *I. gabonensis* kernel oil is mainly composed by saturated fatty acids (SFA) (approximately 95.46%) and, consequently, has a high oxidative stability, even at high temperatures (Yamoneka et al. 2015; Lieb et al. 2018).

Amongst triacylglycerols the most abundant are LaMM (31.1%), CMM/LaLaM (25.6%) and MMM/LaMP (12.9%) (Lieb et al. 2018). Similar results were reported by Meara and Patel (1950), Silou et al. (2011) and Yamoneka et al. (2015). The high proportions of saturated TAGs (approximately 83% tri- and 13% disaturated TAGs), and low proportions of unsaturated TAGs (1.5% di- and 2.6% triunsaturated TAGs), found in *I. gabonensis* kernel fat, could be further associated with the high saturation level of fatty acids previously described (Lieb et al. 2018).

Meara and Patel (1950) described the impossibility of completely separating fully saturated glycerides from mono-oleo-glycerides by crystallization, as well as an individual triglyceride fraction by exhaustive crystallization, in dika fat. The greater solubility of the lower molecular weight, fully

**Table 5.** Micronutrient content (mg/100 g) of *I. gabonensis* kernel.

	Reference		
	Onyeike and Acheru 2002	Ayivor et al. 2011	Dosumu et al. 2012
Iron	0.315	19.374	10.101
Copper	0.139	5.722	2.346
Zinc	0.285	5.786	4.386
Potassium	15.600	0.723	612.55
Sodium	2.020	4.383	59.99

**Table 4.** Physicochemical parameters of *I. gabonensis* kernel oil.

	Reference							
	Onyeike and Acheru 2002	Joseph 1995	Ogunsina et al. 2012	Okoronkwo et al. 2014	Matos et al. 2009	Zoué et al. 2013	Etong et al. 2014	Ifeduba et al. 2013
<b>Chemical parameters</b>								
Oil yield (%)	62.80	66.5	68.37	68.81	62.67	69.76	22.50	64.85
Acid value (mg KOH/g)	24.7	—	—	3.18	12.94	4.67	9.40	—
Saponification value (mg KOH/g)	701	219.2	256.5	230.95	199.50	233.75	187.90	243.90
Peroxide value (meq O <sub>2</sub> /kg fat)	0.04	1.98	0.5	2.67	1.9	3.33	1.80	—
Iodine value (mg I <sub>2</sub> /100 g fat)	21.5	4.2	8.2 g	13.40	4.3	32.43 g	4.50	4.00
Free fatty acids (%)	1.19	0.30	2.72	1.59	4.61	2.33	4.70	—
Unsaponifiable matter (%)	1.70	0.12	—	—	—	1.50	1.50	—
<b>Physical parameters</b>								
State at room temperature	Semi liquid	—	Solid	—	—	—	Solid	—
Specific gravity	0.89	0.85	—	—	—	—	0.88	—
Smoke point (°C)	—	213.0	—	147.57	—	—	78	—
Cloud point (°C)	—	35.0	—	41.83	—	—	—	—
Flash point (°C)	—	—	—	335.33	—	—	—	—
Fire point (°C)	—	—	—	340.67	—	—	—	—
Setting point (°C)	26.3	—	—	—	—	—	25.30	—
Melting/freezing point (°C)	56.0	39.5	—	32.47	—	—	13	—
Color	Golden yellow	—	White	—	—	—	Grey yellow	—

**Table 6.** Mean values (%) of the fatty acid profile of *I. gabonensis* kernel oil.

	Reference								
	Matos et al. 2009	Nangue et al. 2011	Silou et al. 2011	Ogunsina et al. 2012	Zoué et al. 2013	Etong et al. 2014	Yamoneka et al. 2015	Lieb et al. 2018	Ifeduba et al. 2013
Capric acid (C 10:0)	1.34	1.54	0.82	—	0.25	—	1.2	1.3	1.9
Lauric acid (C 12:0)	39.37	40.70	33.18	27.63	39.35	39.40	37.11	37.6	43.3
Myristic acid (C 14:0)	50.92	49.05	55.74	61.68	20.54	20.50	49.83	51.3	45.15
Palmitic acid (C 16:0)	4.97	5.06	5.85	7.49	10.39	10.30	5.51	5.4	5.05
Stearic acid (C 18:0)	0.73	2.38	0.76	0.81	11.46	11.40	0.89	1.0	0.95
Oleic acid (C 18:1)	1.97	0.49	1.35	2.12	6.99	6.90	4.31	2.3	2.95
Linoleic acid (C 18:2)	0.48	—	0.44	0.27	0.01	6.40	1.09	—	0.7
Linolenic acid (C 18:3)	0.00	—	—	—	6.44	—	—	—	—
Arachidic acid (C 20:0)	—	—	—	—	4.52	—	—	—	—
Saturated fatty acid	97.33	98.73	—	97.61	86.56	—	94.54	97.1	96.35
Monosaturated fatty acid	1.97	—	—	2.21	6.99	—	4.31	2.6	2.95
Polyunsaturated fatty acid	0.48	0.49	—	0.27	6.45	—	1.09	0.4	0.7
n-6/n-3 ratio	—	—	—	—	1.55	—	—	—	—

saturated glycerides is more similar to those of the mono-oleoglycerides present than fully saturated components, which explains the problems in separating these fractions. Moreover, the authors noticed that triglycerides occurred in different fractions of the fat, indicating that those are the components primarily responsible for the 'mutual solubility' effects, which precluded the achievement of sharp separations by exhaustive crystallization.

Additionally, unsaponifiable matter composition, as well as phospholipid fraction, have also been discussed. Phospholipid portion (PL) of *I. gabonensis* seed oil has a higher unsaturated fatty acid content when compared with crude seed oil (32.6–36.1% oleic acid, 24.7–29.6% palmitic acid and 16.7–18.0% stearic acid). HPLC analysis of the PL fraction showed that phosphatidylcholine was the most abundant class, making up to 70% of the total PL content followed by phosphatidylinositol and phosphatidylethanolamine (Ifeduba et al. 2013).

Amidst phytosterols identified in *I. gabonensis* seed oil, 4-desmethylsterols were the most representative. More specifically,  $\beta$ -sitosterol and stigmasterol were the predominant components of the unsaponifiable fraction, followed by  $\Delta^5$ -avenasterol and the phytosterol precursor squalene (Silou et al. 2011; Lieb et al. 2018). Lieb et al. (2018) also identified minor 4-desmethylsterols in the seed oil, including cholesterol, 24-methylenecholesterol, campesterol, clerosterol, sitostanol, and  $\Delta^7$ -avenasterol.

Determination of tocochromanols in *I. gabonensis* fat generated contradictory results in literature. While Ifeduba et al. (2013) reported the prevalence of  $\beta$ -tocopherol, followed by  $\alpha$ - and  $\gamma$ -tocopherol, Lieb et al. (2018), contrariwise, identified  $\gamma$ -tocopherol as the most abundant tocopherol homolog and trace amount of  $\alpha$ -tocopherol. Despite the differences, both studies reported similar concentrations of tocopherols in *I. gabonensis* seed oil (4.5–4.6 mg/100 g).

Due to *I. gabonensis* extraordinary proportions of saturated fatty acids and TAGs, it presents a high content of solid fats (SFC) at room temperature, even in tropical countries. Its SFC melting profile remains nearly unchanged up to 20 °C and is completely melted at around 40 °C. After a tempering at 26 °C for 40 h, *I. gabonensis* SFC melting

profile was not affected, indicating that it probably crystallizes directly in a stable form (Yamoneka et al. 2015). Thermograms of *I. gabonensis* fat melting, obtained by differential scanning calorimetry (DSC), also indicate initial melting around 20–25 °C, a fast drop during melting interval with a single maximum peak at 39–40 °C and its termination at 42–45 °C (Yamoneka et al. 2015; Silou et al. 2011; Lieb et al. 2018). As for its polymorphic behavior, *I. gabonensis* seed oil tends to form  $\beta'_1$ -form crystals even after tempering and storage, contrarily to cocoa butter, which presents more stability in  $\beta$ -form. This distinctive polymorphism indicates that tempering would not be necessary to induce conformation stability in *I. gabonensis* seed oil manufacturing (Yamoneka et al. 2015).

Polyphenol composition is described in the following section, as it is related to *I. gabonensis* seed functional properties.

### Functional properties

*In-vivo* and *in-vitro* assays have already been developed to functionally characterize the seed. Data comprising of the antioxidant capacity, total phenol (TPC), total flavonoid (TFC), total anthocyanin (TAC) and total tannin (TTC) contents, as well as total carotenoid (TCC) and ascorbic acid contents, are described in Table 7.

The presence of steroids, flavonoids, alkaloids, cardiac glycosides, volatile oils, terpenoids, tannins and saponins in *I. gabonensis* kernel extract has been revealed on phytochemical screening by Obianime and Uche (2010).

Giami, Okonkwo, and Akusu (1994) studied the influence of heat treatment in the composition of *I. gabonensis* seed flour and stated that increase in temperature occasioned an undesired loss in ascorbic acid, total carotenoid and total polyphenol contents.

*I. gabonensis* seed phytochemical constituents were also compared with mango (*Mangifera indica*) kernels and with a mix of both species (Arogba and Omede 2012; Arogba 2014). According to DPPH, lipid peroxidation and FRAP assays, mango kernels had a higher antioxidant activity than *I. gabonensis* kernels, contrary to nitric oxide assay results. However, *I. gabonensis* kernel results were similar or higher than mango kernel for ascorbic acid content. Total phenol,

**Table 7.** Phytochemical constituents and antioxidant capacity of *I. gabonensis* kernel.

	Value	Reference
DPPH antiradical assay	177.22% (IC <sub>50</sub> )	Arogba and Omede 2012
FRAP assay	431.58 mg of catechin equiv/g 65.43 mM Fe <sup>2+</sup> (IC <sub>50</sub> )	Agbor et al. 2005 Arogba 2014
Lipid peroxidation assay	375.38% (IC <sub>50</sub> )	Arogba 2014
Nitric oxide assay	106.12% (IC <sub>50</sub> )	Arogba 2014
TPC	2.6 mg/100 g 10.74 mg/g 1.15 mg/g dw	Giami et al. 1994 Agbor et al. 2005 Arogba 2014
TFC	0.77 mg QUE/g dw	Arogba 2014
TAC	0.67 ng cyanidin chloride/g dw	Arogba 2014
TTC	1.25 mg catechin/g dw	Arogba 2014
TCC	3.6 mg/100 g	Giami et al. 1994
Ascorbic acid	6.2 mg/100 g	Giami et al. 1994

FRAP, ferric reducing antioxidant power; TPC, total phenolic compounds; TFC, total flavonoid compounds; TAC, total anthocyanin compounds; TTC, total tannin compounds; TCC, total carotenoid content.

flavonoid and tannin contents were also higher in *Mangifera indica* samples, whereas, *I. gabonensis* kernels presented a much higher anthocyanin content. The author showed in these studies that processed kernels of mango (*Mangifera indica*) and *I. gabonensis* contain significant amounts of gal-  
lotannins with high antioxidant capacity even with statistically ( $p < 0.05$ ) higher activity than some other known naturally-occurring phenolic antioxidants (Arogba and Omede 2012; Arogba 2014).

*I. gabonensis* kernel was also compared to 13 Cameroonian herbs/spices. It presented the highest FRAP-free antioxidant capacity followed by *Thymus vulgaris* and ranked third in FRAP total antioxidant but had one of the lowest results in Folin total antioxidant assay (Agbor et al. 2005).

Besides spectrophotometry quantification, it is essential to determine the chemical composition and marker compounds in *I. gabonensis* seed oil for fully comprehend its functionality. Sun and Chen (2012) identified forty-one phenolic compounds in *I. gabonensis* seeds, using an ultrahigh-performance liquid chromatography high-resolution mass spectrometry (UHPLC-HRMS) method. Major constituents in the PDA and total ion chromatograms were identified as ellagic acid and ellagic acid derivatives (di-O-methyl ellagic acid, tri-O-methyl ellagic acid, monomethyl-ellagic acid, di-O-methyl-ellagic acid hexoside, methyl ellagic acid-O-deoxyhexose, di-O-methyl ellagic acid-O-pentoside, di-O-methyl-ellagic acid deoxyhexoside and methyl-ellagic acid-O-rhamnosyl-rhamnoside). Ellagitannins (di-HHDP-hexose and unknown ellagitannins) and flavonoids (kaempferol 3-O-glucoside, quercetin 3-O-rhamnoside, rhamnetin/isorhamnetin with one hexose group and one rhamnosyl-rhamnose and O-methylated flavon) were also identified in minor proportions (Sun and Chen 2012).

Obianime and Uche (2010) studied the effects of *I. gabonensis* seed phytoconstituents in an *in vivo* study, which described the influence of aqueous extract of *I. gabonensis* kernels on biochemical parameters of adult male guinea pigs. The animals were divided into groups in order to perform time-dependent and dose-dependent studies. Groups 1–5 were administered a fixed dose of *I. gabonensis* extract (400 mg/kg/day) over a period of 7, 14, 21, 28 days, respectively. Groups 6–10 were administered different doses of the extract (50–400 mg/kg/day) for 96 hours. Results showed

that the aqueous extract of *I. gabonensis* kernels caused a dose and time-dependent decrease in urea, uric acid, creatine, total cholesterol, protein, alkaline acid, and prostatic phosphatases. Pretreatment with *I. gabonensis* was also able to inhibit the increase in most biochemical parameter levels caused by cadmium administration. The highest reduction effect was obtained with uric acid at 400 mg/kg of *I. gabonensis* extract while the least effect was observed in total cholesterol (Obianime and Uche 2010).

*I. gabonensis* seed extracts were also evaluated for obesity management (Ngondi, Oben, and Minka 2005). The subjects ingested three capsules, three times daily, each containing 350 mg of *I. gabonensis* seed extract (active formulation) or oat bran (placebo), for one month. After 4 weeks, the mean body weight of the *I. gabonensis* group had decreased by 5.26% and that of the placebo group by 1.32%. By the second week, the systolic blood pressure was significantly reduced by the active extract. Obese patients under *I. gabonensis* treatment also had a reduction of 39.21% in total cholesterol, 44.9% in triglycerides, 45.58% in LDL and 32.36% in blood glucose level, as well as an increase of 46.85% in HDL-cholesterol.

Dosumu et al. (2012) studied more specifically the antimicrobial effect of three Nigerian condiments, including *I. gabonensis* dried seed extracts. Clinical isolates of bacteria strains (*Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Salmonella typhi*) and fungi (*Candida albicans*, *Aspergillus niger*, *Rhizopus stolon*, *Penicillium notatum*, *Tricophyton rubrum*, *Epidermophyton floccosum*) were used in the study. Overall, *I. gabonensis* extract had lower antimicrobial activity than the other Nigerian condiments tested. All condiments tested were more effective against fungi than against bacteria. *I. gabonensis* seed extracts obtained with ethyl acetate (200 mg/ml) and methanol (200 mg/ml) presented considerable fungal inhibition when compared to the positive control (Tioconazole, 10 µg/ml).

### Traditional processing and products obtained

The first step in processing *I. gabonensis* kernels is separating the seeds from the mesocarp, using three principal methods for this operation: 'Fresh Cracking'; 'Wet Cracking'



and 'Dry Cracking'. The 'Fresh Cracking' method was reported by Ayuk et al. (1999) and Nkwatoh et al. (2010), in which the whole ripe fruit (pulp and seed) is split in half, through its natural longitudinal line of weakness, with a cutlass or sharp knife. On the other hand, for the other methods, *I. gabonensis* fruits are piled up in heaps and left to fermentate before seed extraction, which facilitates this operation. After fermentation, the seeds can be sun-dried ('Dry Cracking') or directly split ('Wet Cracking'), using truncheons or hard stones as helping tools (Ejiofor, Onwubuke, and Okafor 1987; Ladipo, Fondoun, and Ganga 1996; Ladipo 1999; Ogunsina, Koya, and Adeosun 2008a). As soon as the seeds are cracked, the kernels wrapped in a dark brown testa are exposed and extracted with a knife (Ladipo 1999). The nut cracking process is, therefore, complicated and the dried kernel-in-shell is brittle, resulting in a large percentage of cotyledons being crushed during the process, thereby reducing the market value of the kernels (Ogunsina, Koya, and Adeosun 2008a, 2008b).

Ogunsina, Koya, and Adeosun (2008a, 2008b) investigated fracture behavior of *I. gabonensis* seed in order to provide baseline data for designing an appropriate nutcracker. The physical analysis revealed that minimum toughness was required for nutshell fracture with the small size nuts loaded along the transverse axis. Furthermore, a machine, whose fracture mechanism was based on the deformation characteristics of dried *I. gabonensis* seeds under uni-axial compression, was fabricated. The experimental machine gave 100% cracking efficiency but with 24% kernel breakage in cracking sun-dried *I. gabonensis* seeds with 6.6% moisture content (w.b.). The machine provided a viable and effective technique for safe *I. gabonensis* kernel extraction. Orhevba et al. (2013) also studied physical and mechanical parameters of *I. gabonensis* seed cracking and the influence of moisture content (13.75% and 8.74%). The two moisture content levels were observed to be the range between which *I. gabonensis* kernels can be extracted with least percentage of crushing. Further decrease in the moisture content will make the kernel brittle, while a higher moisture level will make the kernel to stick to the shell, therefore, resulting in crushing during cracking. A motorized machine that is capable of multiple cracking of dika nuts was designed, fabricated and tested by Ogundahunsi, Ogunsina, and Ibrahim (2016). The device utilizes the impact of a sliding hammer block falling from a height to crack a tray of 20 nuts; cracking and splitting them, liberating the embedded kernels as split cotyledons. The highest cracking efficiency and throughput values (72% and 12.86 kg/h, respectively) were obtained for big roasted nuts. The method of pretreatment and dika nut sizes were found to affect the cracking efficiency and throughput of the motorized dika nut cracking machine. After being removed, the kernels are dried for 2 to 7 days, in the sun or on bamboo drying racks over the fireplace (Tchoundjeu, Atangana, and Degrande 2005), in order to remove all moisture (Onimawo et al. 2003; Nkwatoh et al. 2010). This procedure guarantees the quality of the product during storage, by preventing it from discoloring

and from fungal degradation (Ladipo 1999; Ainge and Brown 2001).

Fermentation helps to increase the protein and nitrogen-free extractives of the seeds, as well as to reduce the fat content, which is an advantage if the kernels are consumed integrally (Ekpe, Umoh, and Eka 2007; Ekundayo, Oladipupo, and Ekundayo 2013). Otherwise, if seeds proceed to further processing, the fat loss may be undesired.

At this point, the kernels can be marketed or subjected to further processing. The cotyledons, without the hull, are pounded with a mortar and pestle (Ekpe, Umoh, and Eka 2007). The kernels can also be milled with a grinding machine (Onimawo et al. 2003), which is a more industrial option (Festus and Ibor 2014). The mash, called 'cake', is then molded manually into convenient sizes and shapes, placed in bags or leaves and smoke dried for a few days over a fireplace (Ekpe, Umoh, and Eka 2007; Caspa et al. 2015).

*I. gabonensis* cake can become too slimy over time because of its high fat content; therefore, for an extended shelf-life, defatting is needed (Ainge and Brown 2001; Festus and Ibor 2014). This operation yields, besides crude fat, defatted cake as a product, which, according to Ejiofor, Onwubuke and Okafor (1987), is still acceptable in terms of its color, taste, texture, and drawability after 9 months of storage under ambient conditions, and is more viscous, with greater emulsifying properties than regular flour. The normal flour and defatted flour from *I. gabonensis* kernels are used as ingredients for the popular Ogbono or draw soup which imparts unique flavor, drawability, and thickening properties to the stew (Agbor 1994; Leakey and Newton 1994; Vivien and Faure 1996), and also as 'dika bread' after being baked (Leakey et al. 2005). Leakey et al. (2005) also identified that the food thickening properties of dika nuts were due to two independent and unrelated traits – drawability and viscosity. They also reported considerable tree-to-tree variation in both these traits as well as in protein and oil levels as previously highlighted by Atangana et al. (2001) and Anegbeh et al. (2003) even between trees from the same site, offering opportunities for genetic selection for these traits in new cultivars developed by farmers (Leakey 2014) and improving the existing farmer domestication of these traditionally-important foods (Leakey et al. 2004). Ogbono soup is one of the cheapest, easiest, and fastest Nigerian soups to prepare (Oktay and Sadıkoğlu 2018). Onabanjo and Oguntona (2003) described the following recipe as the most representative of this dish: *I. gabonensis* nuts, bitter (*Vernonia amygdalina*) leaves and okra (*Abelmoschus esculentus* (L.) Moench.) cooked with dried fish, crayfish, ground pepper, pepper, palm oil, bouillon cubes, salt, and water. The influence of *I. gabonensis* seed flour fat content and time of storage on the sensory characteristics of the Ogbono soup was evaluated. Sensory parameters of sliminess (viscosity), taste, aroma, color, and overall acceptability showed that soups prepared from partially defatted *I. gabonensis* seed flour samples (especially the samples with 9% and 12% fat) were more acceptable to the panelists than soups prepared from full-fat *I. gabonensis* seed

flour (Idowu et al. 2013). The preference test carried out by Idowu et al. (2013) during the 12-week period of storage also showed that sliminess, color, taste, aroma, and overall acceptability of Ogbono soups prepared from defatted *I. gabonensis* seed flour (12% and 9% fat) samples packaged in low- and high-density polyethylene films were all acceptable to the panelists. However, the full-fat flour had its sensory parameters significantly decreased in a period of 4 weeks (Akusu and Kiin-Kabari 2013; Idowu et al. 2013).

The defatted cake can also be extruded and molded into Ogbono cubes, which are sold as a convenient cooking ingredient, used as thickeners in Ogbono soup (Okafor, Okolo, and Ejiofor 1996; Ejiofor and Okafor 1997). Bamidele, Ojedokun, and Fasogbon (2015) and Kiin-Kabari and Akusu (2017) developed and analyzed a 'ready-to-cook' powder mix (*I. gabonensis* seed powder, crayfish, stock fish, Ugwu, mixture of locust bean, onion mix, seasoning, and Cameroon powder) for Ogbono soup. Five formulations of instant Ogbono premix were evaluated by Bamidele, Ojedokun, and Fasogbon (2015) (proximate composition, functional properties, micronutrients and sensory analysis). Moisture, protein and carbohydrate contents increased as *I. gabonensis* seed powder percentage decreased in formulations, inversely to fat content. Sensory evaluation showed that the samples with higher percentage of *I. gabonensis* seed powder rated the highest on overall acceptability based on the fact that they showed the real attribute of Ogbono soup that people like which is attributed to the quantity of *I. gabonensis* kernel powder added to the sample (Bamidele, Ojedokun, and Fasogbon 2015). These results are similar to that obtained by Kiin-Kabari and Akusu (2017), which tested formulations of *I. gabonensis* seed (Ogbono) and melon (Egusi) seeds soup premix. They concluded that consumers who do not like very thick soups and low drawability would prefer the formulation with 40:60 Ogbono/Egusi ratio, while consumers who prefer thick soups but low drawability will go for the formulation of 100% 'Egusi'.

Although oil is the major constituent of the seed, according to Nwokocha and Williams (2014), the defatted seed flour essentially consists of polysaccharides with lower than 5% of non-polysaccharide constituents. Nwokocha and Williams (2014) extracted *I. gabonensis* seed gum from its defatted flour by removing soluble sugars and organic pigments with 95% ethanol, followed by dispersion in distilled water (2% w/w), stirring (6 h), and double centrifugation (2500 rpm for 2 h) at 25 °C. On the other hand, Ndjouenkeu et al. (1996) heated the diluted flour (10 g/250 mL) under reflux and then centrifuged the mixture (2000g for 30 min), repeating the process with the supernatant (2 times), then precipitated the crude polysaccharide with 85% ethanol and purified the extract by protein removal. Both studies concluded that *I. gabonensis* seed gum has polyelectrolyte properties, as it is an arabinogalactan but also contains a small proportion of neutral sugars and uronic acids (Ndjouenkeu et al. 1996; Nwokocha and Williams 2014). It showed non-Newtonian behavior at concentrations from 0.2 to 3.0%, having mostly viscous response at concentrations less than

1.0% and elastic response at higher concentrations (Nwokocha and Williams 2014).

Ogaji, Nan, and Hoag (2012) developed a largely physical method for simultaneous extraction of the lipid and polymeric portions of *I. gabonensis*, which was simple, safer, and less expensive than the traditional use of *n*-hexane to extract the lipids. This method was also able to efficiently remove impurities from the gum fraction. The physicochemical properties of the extractives were evaluated, and the results showed similarities in the extractives obtained by this method and those obtained by conventional methods.

Uzomah and Ahiligwo (1999) studied the rheological properties of achi (*Brachystegia eurycoma*) and Ogbono (*I. gabonensis*) seed gums and their potential use as stabilizers in ice cream production. Ogbono seed gum (OSG) cream obtained similar results for quality parameters (maximum overrun, viscosity, shape factor, and meltdown) as the control sample. However, OSG imparted some viscosity to the mixture which resulted in a poor ability to trap and hold air and a poor tendency to resist melting, all of which are characteristics of a satisfactory ice cream. That being said, *I. gabonensis* seed gum was found to be unsuitable as an ice cream stabilizer (Uzomah and Ahiligwo 1999).

It was found that the fat extracted from the kernels can be used for food applications, such as food additive, flavor ingredient, coating fresh citrus fruits and in the manufacture of margarine, oil creams, cooking oil, and defoaming agent. It is also suitable for soap, cosmetics, pharmaceutical products and lather shaving cream (Ejiofor, Onwubuke, and Okafor 1987; Ogunsina et al. 2012; Zoué et al. 2013; Okoronkwo et al. 2014; Etong, Mustapha, and Taleat 2014; Omoniyi et al. 2017).

Matos et al. (2009) characterized margarine made from *I. gabonensis* kernels from two different origins, with and without lecithin. The major fatty acids found in these margarines were oleic acid (35.5%–37%), palmitic acid (18.5%–19.5%) and lauric acid (13.1%–15.1%). The margarines were more unsaturated than the original oil and could be regarded as an oleic acid source. The ratio between linoleic acid (7.07%) and linolenic acid (0.63%) was lower than 2%, showing it can be used for frying food.

*I. gabonensis* seed oil has also been studied as a possible biodiesel source (Bello et al. 2011; Adekunle et al. 2016). It was observed that the kernel fat has similar properties to diesel fuel and superior cold flow properties and flash point, which makes it a suitable alternative fuel for diesel engines (Bello et al. 2011). Adekunle et al. (2016) also concluded that the degumming process improves the physicochemical and biodiesel properties of *I. gabonensis* seed fat, as well as other vegetable oils.

## Other parts

Bark and leaves from *I. gabonensis* have been traditionally used in Nigeria, Cameroon, and other countries where the fruit is available. Proximate composition of stem bark, leaf, and root bark from *I. gabonensis* reveals them to be nutritionally rich (Table 8).

**Table 8.** Proximate composition (%) of *I. gabonensis* stem bark, leaf and root bark (Ezeabara and Ezeani 2016).

	Stem bark	Leaf	Root bark
Moisture	9.43	10.83	8.91
Dry matter	90.58	89.17	91.09
Ash	7.72	9.61	6.58
Fiber	11.38	15.34	8.69
Fat	2.78	1.86	1.45
Protein	5.28	14.78	5.92
Carbohydrates	63.43	47.58	68.44

*I. gabonensis* leaves have been reported to be used as a self-care plant for icterus treatment, by Benin habitants (Allabi et al. 2011). The appropriation of *I. gabonensis* leaves may be associated with high levels of phytochemicals. Ezeabara and Ezeani (2016) reported that *I. gabonensis* leaves contained 2.44% alkaloids, 1.07% flavonoids, and 2.37% anthraquinone, which can be considered high compared to other parts from the same plant. Awah et al. (2012) compared the free radical scavenging activity and phenolic contents from Nigerian medicinal plants and revealed that *I. gabonensis* had high results when compared to the other samples. However, *I. gabonensis* extract presented relative toxicity to humans in the WST-1-based cytotoxicity and cell viability assays (Awah et al. 2012). This toxicity might be related to the high hydrogen cyanide content of 3.45% (Ezeabara and Ezeani 2016).

*I. gabonensis* stem and root barks, instead, do not present relevant levels of hydrogen cyanide, 1.87% and 1.66%, respectively (Ezeabara and Ezeani 2016). Two research studies involving farmers and collectors in Cameroon revealed that *I. gabonensis* stem bark is popular as a traditional medicine. It is reported to treat hernia, yellow fever, dysentery, diarrhea, malaria, to relieve abdominal pain in women, and as antidote for poisons (Ayuk et al. 1999; Zirihi et al. 2005; Caspa et al. 2015).

These effects must be associated with the phytochemicals present in the stem bark. Ezeabara and Ezeani (2016) noted that *I. gabonensis* bark was the most valuable part of the plant, in terms of functional constituents. It consists of the highest percentages of alkaloids (2.78%), flavonoids (1.17%), tannins (1.05%), saponins (0.91%), sterols (0.25%), phenols (0.18%) and anthraquinones (3.17%), compared to the leaf, root bark and raw seed from the same species (Ezeabara and Ezeani 2016).

Zirihi et al. (2005) tested the antiparasmodial activity of ethanol extracts of West African plants, including *I. gabonensis*, and concluded that stem bark extract (10 mg/ml) had a weak antiparasmodial activity against *Plasmodium falciparum* with IC<sub>50</sub> value of 21.6 µg/ml. However, in a study investigating the analgesic effect, the water extract of the stem bark of *I. gabonensis*, when administered to male mice, was found to protect the mice from pain stimuli (Okolo et al. 1995). Another *in vivo* assay evaluated long-term effects of *I. gabonensis* and other two plants, also known to be hypoglycemic, on the oxidative status of normal rabbits (Omonkhua and Onoagbe 2012). Oxidative status was determined by measuring activities of superoxide dismutase (SOD) and catalase (CAT), and the concentration of malondialdehyde (MDA). *I. gabonensis* extract had positive effects

on increasing serum and tissue antioxidant enzymes, particularly in the pancreas, and on decreasing liver MDA levels (Omonkhua and Onoagbe 2012).

## Conclusion

As shown in this work, *I. gabonensis* is a good source of nutrients and phytochemicals and its seeds are already widely consumed and processed traditionally. This review enhances our knowledge about the use of other parts of the fruit, especially the pulp, and about improving the existing methods for a safer and more efficient production of value-added products.

*I. gabonensis* pulp is suitable for juice and wine production, can be also consumed osmose-dried or raw and used as flavourant in the development of other products. It is a vitamin C rich fruit (51–76 mg/100 g) having higher ascorbic acid content than mango or orange. Carotenoids, phenolic compounds, and other phytochemical constituents have also been determined, as well as the hypolipidemic effect of *I. gabonensis* juice administration *in vivo*.

*I. gabonensis* kernel proximate composition revealed its high fat content, as well as a relevant carbohydrate content. *I. gabonensis* kernel oil is considered a 'technical' fat because it resists thermo oxidative, hydrolytic, and enzymatic activities due to its fatty acid profile. *I. gabonensis* seed oil has a unique melting profile and polymorphism, which characterizes it as a possible ingredient for diverse industries, as it remains solid in room temperature (even in tropical countries) and does not require tempering, as it naturally crystallizes in an stable  $\beta_1$ -form. The presence of steroids, flavonoids, alkaloids, cardiac glycosides, volatile oils, terpenoids, tannins and saponins in *I. gabonensis* kernel extract has been revealed on phytochemical screening and its hepatoprotective, nephroprotective, hypolipidemic effects and its influence on body weight have been confirmed. Potential anti-carcinogenic, anti-lipidemic, analgesic and anti-inflammatory effects of the kernel have been highlighted.

Various methods are employed to process the seed. However, all of them have safety issues, when it comes to cracking operation. Various researchers indicated that fermented seeds with specific moisture content, with the appropriate equipment, are easier to crack and kernel loss is reduced. Main products from traditional processing are the sun-dried kernels and the 'cake' which is used as a thickener in 'ogbono soup', a conventional African food. It was described, that with a more sophisticated process, the fat could be extracted from the kernel powder generating a defatted *I. gabonensis* cake. This product can be used as a thickener, stabilizer and as a kind of gum. Not only the defatted flour is potentially manufacturable, but also the crude seed oil, which could be exploited as a potential substitute for fats largely employed, such as cocoa butter and palm oil.

Furthermore, the production of *I. gabonensis* value-added products could reduce food loss, as this would allow the whole fruit to be used. This will also encourage the consumption of wild fruits and support plant biodiversity. This



new approach could ameliorate the diet of rural communities as the *I. gabonensis* fruits are a good source of nutrients and phytochemicals. Commercialization of *I. gabonensis* derived products can also increase the income of the rural communities.

## References

- Achinewhu, S. 1983. Ascorbic acid content of some Nigerian local fruits and vegetables. *Qualitas Plantarum Plant Foods for Human Nutrition* 33 (4):261–6. doi: [10.1007/BF01094751](https://doi.org/10.1007/BF01094751).
- Adekunle, A. S., J. A. O. Oyekunle, O. R. Obisesan, O. S. Ojo, and O. S. Ojo. 2016. Effects of degumming on biodiesel properties of some non-conventional seedoils. *Energy Reports* 2:188–93. doi: [10.1016/j.egy.2016.07.001](https://doi.org/10.1016/j.egy.2016.07.001).
- Agbor, G., J. Oben, J. Ngogang, C. Xinxing, and J. Vinson. 2005. Antioxidant capacity of some herbs/spices from Cameroon: A comparative study of two methods. *Journal of Agricultural and Food Chemistry* 53 (17):6819–24. doi: [10.1021/jf050445c](https://doi.org/10.1021/jf050445c).
- Agbor, L.O.N. 1994. Marketing trends and potentials for *Irvingia gabonensis* products in Nigeria. Paper presented at the ICRAF-IITA Conference on *Irvingia gabonensis*, Ibadan, Nigeria, May.
- Aina, J. 1990. Physico-chemical changes in African mango (*Irvingia gabonensis*) during normal storage ripening. *Food Chemistry* 36 (3): 205–12. doi: [10.1016/0308-8146\(90\)90055-9](https://doi.org/10.1016/0308-8146(90)90055-9).
- Aina, J., and A. Adesina. 1991. Quality attributes of jams from low-usage tropical fruits. *Food Chemistry* 40 (3):345–51. doi: [10.1016/0308-8146\(91\)90118-8](https://doi.org/10.1016/0308-8146(91)90118-8).
- Ainge, L., and N. Brown. 2001. *Irvingia gabonensis* and *Irvingia wombolu*. A state of knowledge. Report undertaken for the Central African Regional Program for the Environment, Oxford Forestry Institute Department of Plant Sciences University of Oxford, UK.
- Akubor, P. 1996. The suitability of African bush mango juice for wine production. *Plant Foods for Human Nutrition (Dordrecht, Netherlands)* 49 (3):213–9.
- Akusu, O. M., and D. B. Kiin-Kabari. 2013. Effect of storage period on selected functional, chemical stability and sensory properties of bush mango (*Irvingia gabonensis*) seed flour. *African Journal of Food Science and Technology* 4 (6):136–40.
- Allabi, A., K. Busia, V. Ekanmian, and F. Bakiono. 2011. The use of medicinal plants in self-care in the Agonlin region of Benin. *Journal of Ethnopharmacology* 133 (1):234–43. doi: [10.1016/j.jep.2010.09.028](https://doi.org/10.1016/j.jep.2010.09.028).
- Amujoyegbe, O., M. Idu, J. Agbedahunsi, and J. Erhabor. 2016. Ethnomedicinal survey of medicinal plants used in the management of sickle cell disorder in Southern Nigeria. *Journal of Ethnopharmacology* 185:347–60. doi: [10.1016/j.jep.2016.03.042](https://doi.org/10.1016/j.jep.2016.03.042).
- Anegbeh, P. O., C. Usoro, V. Ukafor, Z. Tchoundjeu, R. R. B. Leakey, and K. Schreckenber. 2003. Domestication of *Irvingia gabonensis*: 3. Phenotypic variation of fruits and kernels in a Nigerian village. *Agroforestry Systems* 58 (3):213–8.
- Arogba, S. 2014. Phenolics, antiradical assay and cytotoxicity of processed mango (*Mangifera indica*) and bush mango (*Irvingia gabonensis*) kernels. *Nigerian Food Journal* 32 (1):62–72. doi: [10.1016/S0189-7241\(15\)30097-7](https://doi.org/10.1016/S0189-7241(15)30097-7).
- Arogba, S., and A. Omede. 2012. Comparative antioxidant activity of processed mango (*Mangifera indica*) and bush mango (*Irvingia gabonensis*) kernels. *Nigerian Food Journal* 30 (2):17–21. doi: [10.1016/S0189-7241\(15\)30029-1](https://doi.org/10.1016/S0189-7241(15)30029-1).
- Atangana, A. R., Z. Tchoundjeu, J. M. Fondoun, E. Asaah, M. Ndoumbe, and R. R. B. Leakey. 2001. Domestication of *Irvingia gabonensis*: 1. Phenotypic variation in fruits and kernels in two populations from Cameroon. *Agroforestry Systems* 53 (1):55–64.
- Awah, F., P. Uzoegwu, P. Ifeonu, J. Oyugi, J. Rutherford, X. Yao, F. Fehrmann, K. Fowke, and M. Eze. 2012. Free radical scavenging activity, phenolic contents and cytotoxicity of selected Nigerian medicinal plants. *Food Chemistry* 131 (4):1279–86. doi: [10.1016/j.foodchem.2011.09.118](https://doi.org/10.1016/j.foodchem.2011.09.118).
- Aworh, O.C. 2014. *Lesser-known Nigerian fruits and vegetables: Post-harvest handling, utilization and nutritional value*. Ibadan, Nigeria: Ibadan University Press.
- Aworh, O. C. 2015. Promoting food security and enhancing Nigeria's small farmers' income through value-added processing of lesser-known and under-utilized indigenous fruits and vegetables. *Food Research International* 76:986–91.
- Ayivor, J. E., S. Debrah, C. K. Nuviadenu, and A. Forson. 2011. Evaluation of elemental contents of wild mango (*Irvingia gabonensis*) fruit in Ghana. *Advance Journal of Food Science and Technology* 3 (5):381–4.
- Ayuk, E., B. Duguma, S. Franzel, J. Kengue, M. Mollet, T. Tiki-Manga, and P. Zenkeng. 1999. Uses, management and economic potential of *Irvingia gabonensis* in the humid lowlands of Cameroon. *Forest Ecology and Management* 113 (1):1–9. doi: [10.1016/S0378-1127\(98\)00323-5](https://doi.org/10.1016/S0378-1127(98)00323-5).
- Bamidele, O., O. Ojedokun, and B. Fasogbon. 2015. Physico-chemical properties of instant ogbono (*Irvingia gabonensis*) mix powder. *Food Science & Nutrition* 3 (4):313–8. doi: [10.1002/fsn3.220](https://doi.org/10.1002/fsn3.220).
- Bello, E. I., A. O. Fade-Aluko, S. A. Anjorin, and T. S. Mogaji. 2011. Characterization and evaluation of African bush mango nut (dika nut) (*Irvingia gabonensis*) oil biodiesel as alternative fuel for diesel engines. *Journal of Petroleum Technology and Alternative Fuels* 2 (9): 176–80.
- Bvenura, C., and D. Sivakumar. 2017. The role of wild fruits and vegetables in delivering a balanced and healthy diet. *Food Research International (Ottawa, Ont.)* 99 (Pt 1):15–30. doi: [10.1016/j.foodres.2017.06.046](https://doi.org/10.1016/j.foodres.2017.06.046).
- Caspa, R. G., I. R. Tchouamo, J. P. Mate Mweru, J. Mbang Amang, and M. Ngang Ngwa. 2015. The place of *Irvingia gabonensis* in village communities around the Lobeke National Park in Cameroon. *Bois & Forêts des Tropiques* 324 (2):5–17. doi: [10.19182/bft2015.324.a31262](https://doi.org/10.19182/bft2015.324.a31262).
- Codex Alimentarius. 2015. Standard for edible fats and oils not covered by individual standards. CODEX STAN 19-1981. Revision: 1987 and 1999. Amendment: 2009, 2013 and 2015.
- Dosumu, O., O. Oluwaniyi, G. Awolola, and O. Oyediji. 2012. Nutritional composition and antimicrobial properties of three Nigerian condiments. *Nigerian Food Journal* 30 (1):43–52. doi: [10.1016/S0189-7241\(15\)30012-6](https://doi.org/10.1016/S0189-7241(15)30012-6).
- Ejiofor, M.A.N. 1994. Nutritional values of ogbono (*Irvingia gabonensis* var. *excelsa*). Paper presented at the ICRAF-IITA Conference on *Irvingia gabonensis*, Ibadan, Nigeria, May.
- Ejiofor, M. A. N., and J. Okafor. 1997. Prospects for commercial exploitation of Nigerian indigenous trees, vegetables, fruits and seeds through food and industrial products formulation. *International Tree Crops Journal* 9 (2):119–29. doi: [10.1080/01435698.1997.9752967](https://doi.org/10.1080/01435698.1997.9752967).
- Ejiofor, M. A. N., S. Onwubuke, and J. Okafor. 1987. Developing improved methods of processing and utilization of kernels of *Irvingia gabonensis* (var. *Gabonensis* and var. *Excelsa*). *International Tree Crops Journal* 4 (4):283–90. doi: [10.1080/01435698.1987.9752829](https://doi.org/10.1080/01435698.1987.9752829).
- Eka, O. U. 1980. Proximate composition of seeds of bush mango tree and some properties of dika fat. *Nigerian Journal of Nutrition Science* 1:33–36.
- Ekpe, O. O., I. B. Umoh, and O. U. Eka. 2007. Effect of a typical rural processing method on the proximate composition and amino acid profile of bush mango seeds (*Irvingia gabonensis*). *African Journal of Food, Agriculture, Nutrition and Development* 7 (1):1–12.
- Ekundayo, F. O., O. A. Oladipupo, and E. A. Ekundayo. 2013. Studies on the effects of microbial fermentation on bush mango (*Irvingia gabonensis*) seed cotyledons. *African Journal of Microbiological Research* 7 (34):4363–7.
- Elah, E. 2010. Markets and market chain analysis for bush mango (*Irvingia* sp.) in the South West and East regions of Cameroon. PhD diss., University of Buea.
- Emejulu, A. A., C. S. Alisi, E. S. Asiwe, K. M. Iheanacho, and V. A. Onwuliri. 2014. Hypolipidemic effect of *Irvingia gabonensis* fruits juice on sodium fluoride induced dyslipidemia in rats. *African*



- Journal of Biochemistry Research* 8 (8):151–7. doi: [10.5897/AJBR2014.0782](https://doi.org/10.5897/AJBR2014.0782).
- Etebu, E. 2012. Differences in fruit size, postharvest pathology and phytochemicals between *Irvingia gabonensis* and *Irvingia wombolu*. *Sustainable Agriculture Research* 2 (1):52–61. doi: [10.5539/sar.v2n1p52](https://doi.org/10.5539/sar.v2n1p52).
- Etong, D. I., A. O. Mustapha, and A. A. Taleat. 2014. Physicochemical properties and fatty acid composition of dikanut (*Irvingia gabonensis*) seed oil. *Research Journal of Chemical Sciences* 4 (12):70–4.
- Ezeabara, C. A., and D. S. Ezeani. 2016. Comparative study of phytochemical and nutrient contents of various parts of *Irvingia gabonensis* (Aubry-Lecomte ex O' Rorke) Baill. and *Irvingia wombolu* Vermoesen. *Scientia Agriculturae* 14 (3):284–8.
- Ezekwe, C. I., and O. Obidoa. 2001. Biochemical effects of *Vernonia amygdalina* on rat liver microsome. *Nigerian Journal of Biochemistry and Molecular Biology* 16:1745–98.
- Falade, K., and O. C. Aworh. 2004. Adsorption isotherms of osmo-dried African star apple (*Chrysophyllum albidum*) and African mango (*Irvingia gabonensis*) slices. *European Food Research and Technology* 218 (3):278–83. doi: [10.1007/s00217-003-0843-8](https://doi.org/10.1007/s00217-003-0843-8).
- Falade, K., and O. C. Aworh. 2005. Sensory evaluation and consumer acceptance of osmosed and oven-dried African star apple and African mango. *Journal of Food, Agriculture and Environment* 3: 91–6.
- FAO. 1968. Food composition table for use in Africa. Accessed December 29, 2018. <http://www.fao.org/docrep/003/X6877E/X6877E00.htm#TOC>.
- Festus, C., and M. Ibor. 2014. The effect of defatting on the sliminess and shelf-life of Ugiri cotyledons. *Archives of Applied Science Research* 6 (6):1–6.
- Franzel, S., H. Jaenicke, and W. Janssen. 1996. *Choosing the right trees: Setting priorities for multipurpose tree improvement*. ISNAR Research Report 8. The Hague, the Netherlands: International Service for National Agricultural Research.
- Fungo, R., J. Muyonga, M. Kabahenda, C. Okia, and L. Snook. 2016. Factors influencing consumption of nutrient rich forest foods in rural Cameroon. *Appetite* 97:176–84. doi: [10.1016/j.appet.2015.12.005](https://doi.org/10.1016/j.appet.2015.12.005).
- Giami, S., V. Okonkwo, and M. Akusu. 1994. Chemical composition and functional properties of raw, heat-treated and partially proteolysed wild mango (*Irvingia gabonensis*) seed flour. *Food Chemistry* 49 (3):237–43. doi: [10.1016/0308-8146\(94\)90166-X](https://doi.org/10.1016/0308-8146(94)90166-X).
- Gopalan, C., B.V. Rama Sari, and S.C. Balasubramanian. 2014. *Nutritive value of Indian foods*. Hyderabad, India: National Institute of Nutrition, Indian Council of Medical Research.
- Harris, D. J. 1996. A revision of the *Irvingiaceae* in Africa. *Bulletin du Jardin Botanique National de Belgique / Bulletin Van de National Plantentuin Van België* 65 (1/2):143–96. doi: [10.2307/3668184](https://doi.org/10.2307/3668184).
- Ibezim, G. N. U. 2015. Chemical constituents of dikanut and soyabean seeds. *Time Journals of Medicinal Plant Sciences and Pharmacology* 3 (2):17–22.
- Idowu, M., S. Omoniyi, F. Henshaw, and I. Olayiwola. 2013. Sensory acceptability of partially defatted dikanut (*Irvingia gabonensis*) flour in ogbono soup. *Journal of Culinary Science & Technology* 11 (4): 346–55. doi: [10.1080/15428052.2013.798607](https://doi.org/10.1080/15428052.2013.798607).
- Ifeduba, E. A., M. N. Awachie, J. S. M. Sabir, and C. C. Akoh. 2013. Fatty acid composition of *Irvingia gabonensis* and *Treculia Africa* seed lipid and phospholipids. *Journal of the American Oil Chemists' Society* 90 (4):517–28. doi: [10.1007/s11746-012-2199-3](https://doi.org/10.1007/s11746-012-2199-3).
- Joseph, J. 1995. Physico-chemical attributes of wild mango (*Irvingia gabonensis*) seeds. *Bioresource Technology* 53 (2):179–81. doi: [10.1016/0960-8524\(95\)00044-F](https://doi.org/10.1016/0960-8524(95)00044-F).
- Joseph, K., and O. Aworh. 1991. Composition, sensory quality and respiration during ripening and storage of edible wild mango (*Irvingia gabonensis*). *International Journal of Food Science & Technology* 26 (3):337–42. doi: [10.1111/j.1365-2621.1991.tb01170.x](https://doi.org/10.1111/j.1365-2621.1991.tb01170.x).
- Joseph, K., and O. Aworh. 1992. Post-harvest treatment of wild mango (*Irvingia gabonensis*) for improved shelf life. *Food Chemistry* 44 (1): 45–8. doi: [10.1016/0308-8146\(92\)90256-2](https://doi.org/10.1016/0308-8146(92)90256-2).
- Kiin-Kabari, D. B., and O. M. Akusu. 2017. Production, proximate, functional and organoleptic assessment of ready-to-cook “ogbono”/”egusi” seeds premix (dry mix powder). *International Journal of Food Science and Nutrition* 2 (1):82–5.
- Kuete, V., and T. Efferth. 2011. Pharmacogenomics of Cameroonian traditional herbal medicine for cancer therapy. *Journal of Ethnopharmacology* 137 (1):752–66. doi: [10.1016/j.jep.2011.06.035](https://doi.org/10.1016/j.jep.2011.06.035).
- Ladipo, D.O. 1999. The development of quality control standard for Ogbono (*Irvingia gabonensis* and *Irvingia wombolu*) kernels: Efforts toward encouraging organized and further international trade in NWFP in West and central Africa. In *Non wood Forest products of Central Africa – Current research issues and prospects for conservation and development*, eds. T. C. H Sundarland, L. E. Clark, and P. Vantomme, 245–50. Rome, Italy: FAO.
- Ladipo, D. O., J. M. Fondoun, and N. Ganga. 1996. Domestication of the bush mango (*Irvingia* spp): Some exploitable intra-specific variation in West and Central Africa. In *Domestication and commercialization of non-timber forest products in agroforestry systems*, eds. R. R. B. Leakey, A. B. Temu, M. Melynyk, and P. Vantomme, 193–205. Rome, Italy: FAO.
- Leakey, R.R.B. 2014. Agroforestry – Participatory domestication of trees. In *Encyclopedia of agriculture and food systems*, ed. N. van Alfen, Vol. 1, 253–69. San Diego, CA: Elsevier Publishers.
- Leakey, R. R. B. 2017. Socially modified organisms in multifunctional agriculture – Addressing the needs of smallholder farmers in Africa. *Archives of Crop Science* 1:20–9.
- Leakey, R. R. B. 2018. Converting ‘trade-offs’ to ‘trade-ons’ for greatly enhanced food security in Africa: Multiple environmental, economic and social benefits from ‘socially modified crops. *Food Security* 10 (3):505–24. doi: [10.1007/s12571-018-0796-1](https://doi.org/10.1007/s12571-018-0796-1).
- Leakey, R. R. B. 2019. From ethnobotany to mainstream agriculture – Socially-modified Cinderella species capturing ‘trade-ons’ for ‘land maxing. *Planta (Special Issue on Orphan Crops)*. doi: [10.1007/s00425-019-03128-z](https://doi.org/10.1007/s00425-019-03128-z).
- Leakey, R. R. B., J. Fondoun, A. Atangana, and Z. Tchoundjeu. 2000. Quantitative descriptors of variation in the fruits and seeds of *Irvingia gabonensis*. *Agroforestry Systems* 50 (1):47–58.
- Leakey, R. R. B., P. Greenwell, M. Hall, A. Atangana, C. Usoro, P. Anegebeh, J. Fondoun, and Z. Tchoundjeu. 2005. Domestication of *Irvingia gabonensis*: 4. Tree-to-tree variation in food-thickening properties and in fat and protein contents of dika nut. *Food Chemistry* 90 (3):365–78. doi: [10.1016/j.foodchem.2004.04.012](https://doi.org/10.1016/j.foodchem.2004.04.012).
- Leakey, R.R.B., and A.C. Newton. 1994. Domestication of ‘Cinderella’ species as the start of a woody-plant revolution. In *Tropical trees: The potential for domestication and the rebuilding of forest resources*, eds. R. R. B. Leakey and A. C. Newton, 3–6. London, UK: HMSO.
- Leakey, R. R. B., and T. Page. 2006. The ‘ideotype concept’ and its application to the selection of cultivars of trees providing agroforestry tree products. *Forests, Trees and Livelihoods* 16 (1):5–16. doi: [10.1080/14728028.2006.9752542](https://doi.org/10.1080/14728028.2006.9752542).
- Leakey, R. R. B., Z. Tchoundjeu, R. I. Smith, R. C. Munro, J.-M. Fondoun, J. Kengue, P. O. Anegebeh, A. R. Atangana, A. N. Waruhiu, E. Asaah, et al. 2004. Evidence that subsistence farmers have domesticated indigenous fruits (*Dacryodes edulis* and *Irvingia gabonensis*) in Cameroon and Nigeria. *Agroforestry Systems* 60 (2): 101–11. doi: [10.1023/B:AGFO.0000013259.95628.22](https://doi.org/10.1023/B:AGFO.0000013259.95628.22).
- Leakey, R.R.B., J.C. Weber, T. Page, J.P. Cornelius, F.K. Akinnifesi, J.M. Roshetko, Z. Tchoundjeu, and R. Jamnadass. 2012. Tree domestication in agroforestry: Progress in the second decade. In *Agroforestry – The future of global land use*, eds. P. K. Nair and D. Garrity, 145–73. New York, NY: Springer.
- Lieb, V. M., L. K. Schuster, H. G. Schmarr, R. Carle, and C. B. Steingass. 2018. Lipophilic compounds and thermal behaviour of African mango (*Irvingia gabonensis* (Aubry-Lecomte ex O'Rorke) Baill.) kernel fat. *International Journal of Food Science & Technology* 54:626–33. doi: [10.1111/ijfs.13940](https://doi.org/10.1111/ijfs.13940).
- Matos, L., J. Nzikou, E. Matouba, V. Pandzou-Yembe, T. Guembot Mapepoulou, M. Linder, and S. Desobry. 2009. Studies of *Irvingia gabonensis* seed kernels: Oil technological applications. *Pakistan Journal of Nutrition* 8 (2):151–7. doi: [10.3923/pjn.2009.151.157](https://doi.org/10.3923/pjn.2009.151.157).

- Mbaeyi, I. E., and L. N. Anyanwu. 2010. Production and evaluation of yoghurt flavoured with solar-dried bush mango (*Irvingia gabonensis*) pulp. *Agro-Science* 9 (2):137–46. doi: [10.4314/as.v9i2.64808](https://doi.org/10.4314/as.v9i2.64808).
- Meara, M., and C. Patel. 1950. The component acids and glycerides of dika fat. *Journal of the Science of Food and Agriculture* 1 (2):48–51. doi: [10.1002/jsfa.2740010205](https://doi.org/10.1002/jsfa.2740010205).
- Nangue, T. J., H. M. Womeni, F. T. Mbiapo, J. Fanni, and L. Michel. 2011. *Irvingia gabonensis* fat: Nutritional properties and effect of increasing amounts on the growth and lipid metabolism of young rats wistar sp. *Lipids in Health and Disease* 10 (1):43. doi: [10.1186/1476-511X-10-43](https://doi.org/10.1186/1476-511X-10-43).
- National Research Council. 2006. *Lost crops of Africa. Volume II: Vegetables*. Washington, DC: The National Academies Press.
- Ndjouenkeu, R., F. Goycoolea, E. Morrisa, and J. Akingbala. 1996. Rheology of okra (*Hibiscus esculentus* L.) and dika nut (*Irvingia gabonensis*) polysaccharides. *Carbohydrate Polymers* 29 (3):263–9. doi: [10.1016/0144-8617\(96\)00016-1](https://doi.org/10.1016/0144-8617(96)00016-1).
- Ngondi, J. L., J. E. Oben, and S. R. Minka. 2005. The effect of *Irvingia gabonensis* seeds on body weight and blood lipids of obese subjects in Cameroon. *Lipids in Health and Disease* 4 (1):12.
- Nkwatoh, A., P. Labode, S. Iyassa, and F. Nkwatoh. 2010. Evaluation of *Irvingia* (*I. gabonensis* and *I. wombulu*) trade; a major non-timber Forest product in the Ejagham Forest Reserve of South West Cameroon. *Ethiopian Journal of Environmental Studies and Management* 3 (1):70–7. doi: [10.4314/ejesm.v3i1.54403](https://doi.org/10.4314/ejesm.v3i1.54403).
- Nwokocha, L., and P. Williams. 2014. Hydrodynamic and rheological properties of *Irvingia gabonensis* gum. *Carbohydrate Polymers* 114: 352–6. doi: [10.1016/j.carbpol.2014.07.071](https://doi.org/10.1016/j.carbpol.2014.07.071).
- Obianime, A., and F. Uche. 2010. The phytoconstituents and the comparative effects of aqueous extract of *Irvingia gabonensis* seeds and proviron on the biochemical parameters of male guinea pigs. *Asian Pacific Journal of Tropical Medicine* 3 (2):101–4. doi: [10.1016/S1995-7645\(10\)60044-2](https://doi.org/10.1016/S1995-7645(10)60044-2).
- Ofundem, T., N. R. Ndiop, A. Abdon, and L. Patrice. 2017. Bush mango (*Irvingia* spp.): Forest and on-farm resource availability and market chains in the Southwest region of Cameroon. *Forests, Trees and Livelihoods* 26 (3):170–82. doi: [10.1080/14728028.2017.1283250](https://doi.org/10.1080/14728028.2017.1283250).
- Ogaji, I., A. Nan, and S. Hoag. 2012. A novel extraction method and some physicochemical properties of extractives of *Irvingia gabonensis* seeds. *Journal of Young Pharmacists* 4 (2):66–72. doi: [10.4103/0975-1483.96612](https://doi.org/10.4103/0975-1483.96612).
- Ogundahunsi, O. E., B. S. Ogunsina, and R. O. Ibrahim. 2016. A motorized device for cracking pre-treated dika nuts. *Agricultural Engineering International: CIGR Journal* 18 (2):309–22.
- Ogunsina, B. S., O. A. Koya, and O. O. Adeosun. 2008a. Deformation and fracture of dika nut (*Irvingia gabonensis*) under uniaxial compressive load. *International Agrophysics* 22:249–53.
- Ogunsina, B. S., O. A. Koya, and O. O. Adeosun. 2008b. A table mounted device for cracking dika nut (*Irvingia gabonensis*). *Agricultural Engineering International: CIGR Journal*. Manuscript PM 08 011, Vol. X.
- Ogunsina, B. S., A. S. Bhatnagar, T. N. Indira, and C. Radha. 2012. The proximate composition of African bush mango kernels (*Irvingia gabonensis*) and characteristics of its oil. *Ife Journal of Science* 14 (1):177–83.
- Ogunsina, B., G. Olatunde, and O. Adeleye. 2014. Effect of pre-treatments on mechanical oil expression from dika kernels. *Nigerian Food Journal* 32 (1):1–9. doi: [10.1016/S0189-7241\(15\)30090-4](https://doi.org/10.1016/S0189-7241(15)30090-4).
- Okafor, J.C., H.C. Okolo, and M.A.N. Ejiofor. 1996. Strategies for enhancement of utilization potential of edible woody forest species of South-Eastern Nigeria. In *The biodiversity of African plants*, eds. L. J. G. van der Maesen, X. M. van der Burgt, and J. M. van Medenbach de Rooy, 684–95. Dordrecht, the Netherlands: Springer.
- Okolo, C. O., P. B. Johnson, E. M. Abdurahman, I. Abdu-Aguye, and I. M. Hussaini. 1995. Analgesic effect of *Irvingia gabonensis* stem bark extract. *Journal of Ethnopharmacology* 45 (2):125–9. doi: [10.1016/0378-8741\(94\)01199-A](https://doi.org/10.1016/0378-8741(94)01199-A).
- Okolo, H.C. 1994. Industrial potential of various *Irvingia gabonensis* products such as oil, Ogbono and juice. Paper presented at the ICRAF-IITA Conference on *Irvingia gabonensis*, Ibadan, Nigeria, May.
- Okoronkwo, C. U., E. E. C. Agoha, O. Alloysius, A. C. Ogodo, and N. O. Nwachukwu. 2014. Physical and chemical characteristics of the African bush mango (*Irvingia gabonensis* var. *garbonensis*) seed oil. *International Journal of Advances in Engineering and Management* 1 (6):28–31.
- Oktay, S., and S. Sadıkoğlu. 2018. The gastronomic cultures' impact on the African cuisine. *Journal of Ethnic Foods* 5 (2):140–6. doi: [10.1016/j.jef.2018.02.005](https://doi.org/10.1016/j.jef.2018.02.005).
- Olayiwola, I., V. Akinfenwa, C. Oguntona, S. Sanni, O. Onabanjo, and W. Afolabi. 2013. Phytonutrient, antioxidant and mineral composition of some wild fruits in South West Nigeria. *Nigerian Food Journal* 31 (2):33–40. doi: [10.1016/S0189-7241\(15\)30074-6](https://doi.org/10.1016/S0189-7241(15)30074-6).
- Omoniyi, S., M. Idowu, A. Adeola, and A. Folorunso. 2017. Chemical composition and industrial benefits of dikanut (*Irvingia gabonensis*) kernel oil. *Nutrition & Food Science* 47 (5):741–51. doi: [10.1108/NFS-03-2017-0054](https://doi.org/10.1108/NFS-03-2017-0054).
- Omonkhua, A. A., and I. O. Onoagbe. 2012. Long-term effects of three hypoglycaemic plants (*Irvingia gabonensis*, *Urena lobata* and *Carica papaya*) on the oxidative status of normal rabbits. *Biokemistri* 24 (2):82–9.
- Onabanjo, O. O., and C. R. B. Oguntona. 2003. Iron, zinc, copper and phytate content of standardized Nigerian dishes. *Journal of Food Composition and Analysis* 16 (6):669–76. doi: [10.1016/S0889-1575\(03\)00063-2](https://doi.org/10.1016/S0889-1575(03)00063-2).
- Onimawo, I., F. Oteno, G. Orokp, and P. Akubor. 2003. Physicochemical and nutrient evaluation of African bush mango (*Irvingia gabonensis*) seeds and pulp. *Plant Foods for Human Nutrition* 58 (3):1–6. doi: [10.1023/B:QUAL.0000040320.33549.df](https://doi.org/10.1023/B:QUAL.0000040320.33549.df).
- Onyeike, E., and G. Acheru. 2002. Chemical composition of selected Nigerian oil seeds and physicochemical properties of the oil extracts. *Food Chemistry* 77 (4):431–7. doi: [10.1016/S0308-8146\(01\)00377-6](https://doi.org/10.1016/S0308-8146(01)00377-6).
- Orhevb, B. A., P. A. Idah, S. E. Adebayo, and C. C. Nwankwo. 2013. Determination of some engineering properties of dika nut (*Irvingia gabonensis*) at two moisture content levels as relevant to its processing. *International Journal of Engineering Research and Applications* 3 (2):182–8.
- Ruth Charrondière, U., B. Stadlmayr, D. Rittenschober, B. Mouille, E. Nilsson, E. Medhammar, T. Olango, S. Eisenwagen, D. Persijn, K. Ebanks, et al. 2013. FAO/INFOODS food composition database for biodiversity. *Food Chemistry* 140 (3):408–12. doi: [10.1016/j.foodchem.2012.08.049](https://doi.org/10.1016/j.foodchem.2012.08.049).
- Shaheen, S., M. Ahmad, and N. Haroon. 2017. Edible wild plants: A solution to overcome food insecurity. In *Edible wild plants: An alternative approach to food security*, eds. S. Shaheen, M. Ahmad, and N. Haroon, 1st ed., 41–57. Heidelberg, Germany: Springer.
- Silou, T., B. W. Loumouamou, D. Nsikabaka, T. Kinkela, M. Nzikou, P. Chalard, and G. Figueredo. 2011. Contribution to the varietal delimitation of *Irvingia gabonensis* and *Irvingia wombulu*. Chemical composition variability of fats extracted from kernels. *Journal of Food Technology* 9 (1):36–42. doi: [10.3923/jftech.2011.36.42](https://doi.org/10.3923/jftech.2011.36.42).
- Singh, A. K. 2007. Bush mango (*Irvingia gabonensis*): New potential multipurpose fruit tree for India. *Indian Journal of Plant Genetic Resources* 20 (1):32–7.
- Stadlmayr, B., U. Charrondière, S. Eisenwagen, R. Jamnadass, and K. Kehlenbeck. 2013. Nutrient composition of selected indigenous fruits from Sub-Saharan Africa. *Journal of the Science of Food and Agriculture* 93 (11):2627–36. doi: [10.1002/jsfa.6196](https://doi.org/10.1002/jsfa.6196).
- Sun, J., and P. Chen. 2012. Ultra high-performance liquid chromatography with high-resolution mass spectrometry analysis of African mango (*Irvingia gabonensis*) seeds, extract, and related dietary supplements. *Journal of Agricultural and Food Chemistry* 60 (35): 8703–9. doi: [10.1021/jf302703u](https://doi.org/10.1021/jf302703u).
- Tchoundjeu, Z., A. Atangana, and A. Degrande. 2005. Indigenous methods in preserving bush mango kernels in Cameroon. *American Journal of Applied Sciences* 2 (9):1337–42. doi: [10.3844/ajassp.2005.1337.1342](https://doi.org/10.3844/ajassp.2005.1337.1342).
- USDA. 2018. Food Composition Databases. Accessed December 29, 2018. <https://ndb.nal.usda.gov/ndb/>.

- Uzomah, A., and R. Ahiligwo. 1999. Studies on the rheological properties and functional potentials of achi (*Brachystegia eurycoma*) and ogbono (*Irvingia gabonensis*) seed gums. *Food Chemistry* 67 (3): 217–22. doi: [10.1016/S0308-8146\(98\)00256-8](https://doi.org/10.1016/S0308-8146(98)00256-8).
- Vihotogbé, R., R. G. van den Berg, and M. S. M. Sosef. 2013. Morphological characterization of African bush mango trees (*Irvingia* species) in West Africa. *Genetic Resources and Crop Evolution* 60 (4):1597–614. doi: [10.1007/s10722-013-9969-0](https://doi.org/10.1007/s10722-013-9969-0).
- Vivien, J., and J.J. Faure. 1996. *Fruitiers sauvages d'Afrique. Espèces du Cameroun*. Clohars Carnoetn, France: Nguila-Kerou Editions.
- Yamoneka, J., P. Malumba, C. Blecker, M. Gindo, G. Richard, M. L. Fauconnier, G. Lognay, and S. Danthine. 2015. Physicochemical properties and thermal behaviour of African wild mango (*Irvingia gabonensis*) seed fat. *LWT - Food Science and Technology* 64 (2): 989–96. doi: [10.1016/j.lwt.2015.06.035](https://doi.org/10.1016/j.lwt.2015.06.035).
- Zirihi, G., L. Mambu, F. Guédé-Guina, B. Bodo, and P. Grellier. 2005. In vitro antiplasmodial activity and cytotoxicity of 33 West African plants used for treatment of malaria. *Journal of Ethnopharmacology* 98 (3):281–5. doi: [10.1016/j.jep.2005.01.004](https://doi.org/10.1016/j.jep.2005.01.004).
- Zoué, L., M. Bédikou, B. Faulet, J. Gonnety, and S. Niamké. 2013. Characterisation of a highly saturated *Irvingia gabonensis* seed kernel oil with unusual linolenic acid content. *Food Science and Technology International* 19 (1):79–87. doi: [10.1177/1082013212442190](https://doi.org/10.1177/1082013212442190).